

# BALKANS PROJECT

## Fieldwork

Initiated in number of licenses

Geochemical sampling -15

Geological mapping-15

Gravity -14


Magneto tellurics -3

**Core drilling -6**


Downgraded/dropped-4

As of March 1<sup>st</sup>, 2013

## ISSUED LICENSES AND WORK PROGRESS




**15 applications in Serbia approved, and first stage exploration completed in all licences**




## BALKANS PROJECT

### LOPARE GRAVITY & SURFACE GEOCHEMISTRY DATA




- An ~100m thick sequence of dolomitic sediments with zones of borate and volcanic ash falls is exposed along southern portion of the license.
- Borate mineral occurs in the form of searlesite ( $\text{NaBSi}_5\text{O}_5(\text{OH})_2$ ) up to 9.8%  $\text{B}_2\text{O}_3$ .**
- Rio Tinto drilled 5 shallow holes (average 303m depth) in a blind test for trona in 2002 without benefit of gravity geophysics, a similar situation to Valjevo in Serbia.
- 2 current holes to drill test entire basin stratigraphy.**




## LOPARE –VEGETATION ANOMALY AND OUTCROP OF BORATE MINERALIZATION WITH GRADES UP TO 9.98% $B_2O_3$





# BALKANS PROJECT

## VALJEVO GRAVITY DATA AND DRILL RESULTS TO DATE



- **World class** (in size) borate occurrence being outlined.
- Only proterite intersected to date indicating on **edge of large system**.
- **Preserved borates over 6km<sup>2</sup>** with pseudomorphs of borates over extra 10km<sup>2</sup> (total 16km<sup>2</sup>).
- Thickness of preserved borate zone ranges from 1.3m to 16.3m.
- **Thickness** of total borate zone (incl. pseudomorphs) **from 31.7m to 67.0m**.
- Highest grades (8% c/o) from holes **VBN-4 and VBN-8** incl. 1.6M @ 18.2% B<sub>2</sub>O<sub>3</sub> and 1.0m @ 17.7% B<sub>2</sub>O<sub>3</sub>.
- High Li<sub>2</sub>O grades up to 18.8m @ 0.26%.
- **Initial focus of 2013 drilling to north east of VBN-4.**
-